



Autologous vs. Non-Myeloablative Allogeneic Hematopoietic Stem Cell Transplantation (HSCT) for Patients with Relapsed Follicular Non-Hodgkin's Lymphoma

BMT CTN PROTOCOL 0202 VERSION 4.0

Core Study Participants:

Case Western Reserve Consortia
Oregon Health & Science University
University Hospitals of Cleveland
Washington University
City of Hope National Medical Center
Dana Farber/Partners Cancer Center
Duke University Medical Center-Adult
Stanford Hospital and Clinics
UCSD/SCRIPPS School of Medicine
University of Florida College of Medicine (Shands)
University of Michigan Medical Center
University of Minnesota
University of Nebraska Medical Center
University of Pennsylvania Cancer Center
University of Texas, MD Anderson Cancer Center

Non-Core Study Participants:

Baylor University Medical Center
Blood and Marrow Transplant Group of Georgia
Emory University
Hackensack University Medical Center
H. Lee Moffitt Cancer Center
Indiana BMT at Beech Grove
Kansas City Blood & Marrow Transplant Center
Karmanos Cancer Institute, BMT
Loyola University Medical Center
Medical College of Wisconsin
Providence Portland Medical Center
Temple University
University of Maryland
University of Pittsburgh
University of Wisconsin
Vanderbilt University Medical Center
Virginia Commonwealth University Medical Center